

HOW TO USE

 **Targretin**[®]
(bexarotene) 75 mg capsules

*and make the most out of
your CTCL treatment*

TARGRETIN Capsules should not be used by women who are pregnant, believe they may be pregnant, or are planning to become pregnant. Please see Important Safety Information Section, including Pregnancy Boxed Warning located on the inside back panel, and Patient Instructions for Use located within the accompanying Prescribing Information.

ELIGIBLE
COMMERCIALLY
INSURED PATIENTS

\$ 0* CO-PAY



* This offer is only valid for eligible patients with commercial insurance. Commercially insured patients whose insurance does not cover TARGRETIN and cash paying patients will pay more. This offer is not valid for prescriptions reimbursed in whole or in part by Medicaid, Medicare, federal or state programs (including any state prescription drug programs). Void where prohibited by law, taxed or restricted. Cannot be redeemed at government-subsidized clinics. Valeant Pharmaceuticals reserves the right to rescind, revoke, or amend this offer without notice.

GETTING THE MOST
OUT OF **TARGRETIN**
CAPSULES TREATMENT



*It's important to take
TARGRETIN Capsules as
your doctor prescribes to get
the most out of treatment. This
includes how many capsules
you should take.*

All medications have side effects. You should call your physician regarding any questions or concerns you may have when taking TARGRETIN Capsules.

THE BASICS ABOUT **TARGRETIN** CAPSULES

TARGRETIN (bexarotene) Capsules, 75 mg are used to treat the skin problems arising from a disease called cutaneous T-cell lymphoma, or CTCL. Your doctor must supervise the use of TARGRETIN Capsules.¹

THERE ARE SOME IMPORTANT THINGS TO KNOW BEFORE STARTING TARGRETIN CAPSULES TREATMENT:¹

- **TARGRETIN Capsules can cause major damage to a fetus.** If you are capable of becoming pregnant, you should take contraceptive measures to avoid pregnancy before, during and after treatment. It is recommended that two reliable forms of contraception be used together.
- TARGRETIN Capsules can greatly increase blood levels of lipids (triglycerides and cholesterol) and these levels must be monitored and, if elevated, treated.
- TARGRETIN Capsules can cause an underactive thyroid and periodic blood tests will be needed to detect this. Medication to control the condition may be necessary.
- Do not take more than the recommended daily dietary allowance of Vitamin A (4000 to 5000 International Units).
- Your skin may become more sensitive to sunlight while using these medicines. Minimize exposure to sunlight and do not use a sunlamp.

You should also tell your doctor or nurse about any current medications or health conditions.

Please see Important Safety Information, including Pregnancy Boxed Warning located on the inside back panel, and Patient Instructions for Use located within the accompanying Prescribing Information.

HOW TO TAKE TARGRETIN (bexarotene) CAPSULES, 75 MG

TARGRETIN Capsules should be taken once a day with a meal. Your doctor will tell you how many you need to take. Take them all at once, swallowing each capsule whole. If you miss a dose, take it as soon as possible. If it's almost time for your next dose, skip the missed dose.¹

If you take too many TARGRETIN Capsules or someone else accidentally takes your medicine, contact your doctor, emergency room or the nearest hospital immediately.


HOW LONG BEFORE YOU CAN EXPECT YOUR CTCL TO IMPROVE ON TARGRETIN CAPSULE TREATMENT?

It is important to use as directed by your physician as a response may take several weeks to several months. Your doctor should determine how long you should be taking TARGRETIN Capsules, and when treatment may be stopped.

SUPPORTING YOU, **ONE DAY AT A TIME**

LIVING WITH CTCL

Living with CTCL is a daily challenge. There will be up days and down days. What's important is having support you can lean on, including your loved ones and health care team. *Valeant also wants to do its part.*

 **Targretin**[®]
(bexarotene) 75 mg capsules

Please see Important Safety Information, including Pregnancy Boxed Warning located on the inside back panel, and Patient Instructions for Use located within the accompanying Prescribing Information.



ENHANCED ACCESS AND SUPPORT DURING TREATMENT WITH **TARGRETIN** CAPSULES:



SUPPORTING YOU, ONE DAY AT A TIME

Living with CTCL is a daily challenge. There will be up days and down days. What's important is having support you can lean on, including your loved ones and health care team. *Valeant also wants to do its part.*

That's why we created the **Targretin Patient Care Program**. It represents our commitment to providing you with the support you need. This program is designed to help you gain access to treatment and provide support throughout the course of your care.

THIS SUPPORT PROGRAM PROVIDES YOU:

→ Customer Service Representatives

To help you have a positive experience when filling your prescriptions.

→ Patient Access Coordinators

To assist you with insurance coverage and help you seek financial assistance, when needed.

→ Pharmacy Technicians

To make sure your prescriptions are processed with the utmost accuracy.

→ Pharmacists

To help you better understand and follow your treatment plan.



TARGRETIN (bexarotene) CAPSULES, 75 MG is a medicine used to treat the skin problems arising from a disease called cutaneous T-cell lymphoma, or CTCL. Your doctor must supervise the use of TARGRETIN Capsules.¹

Before starting treatment with TARGRETIN Capsules, make sure you tell your doctor about all your medical conditions and any medications you are currently taking.

Please see **Important Safety Information Section, including Pregnancy Boxed Warning located on the inside back panel, and Patient Instructions for Use located within the accompanying Prescribing Information.**



REFERENCE:

1. TARGRETIN® (bexarotene) capsules, 75 mg Prescribing Information. 2013.

®/TMs are trademarks of Valeant Pharmaceuticals International, Inc. or its affiliates. Any other product or brand names and logos are the property of their respective owners.

©2015 Valeant Pharmaceuticals North America LLC.

DM/TGC/15/0001

INDICATION

TARGRETIN® (bexarotene) Capsules, 75 mg are used to treat the skin problems arising from a disease called cutaneous T-cell lymphoma, or CTCL, in patients who have not responded well to other treatments. Your healthcare provider must supervise the use of TARGRETIN Capsules.

IMPORTANT SAFETY INFORMATION INCLUDING PREGNANCY BOXED WARNING

What are the most important things I should know about TARGRETIN Capsules?

Do not take TARGRETIN Capsules if you are pregnant or if you plan to become pregnant.

- ➔ TARGRETIN Capsules may harm your fetus (unborn baby). You should contact your healthcare provider immediately if you believe or suspect you are pregnant while you are taking TARGRETIN Capsules and until one month after you stop taking TARGRETIN Capsules.
- ➔ If you are capable of becoming pregnant, you must have a pregnancy test, within one week before you start TARGRETIN Capsule therapy and monthly while you are taking TARGRETIN Capsules, confirming you are not pregnant.
- ➔ You must use effective contraception (birth control) continuously starting one month before beginning treatment with TARGRETIN Capsules until one month after you stop taking TARGRETIN Capsules. It is strongly recommended that two reliable forms of contraception be used together. At least one of these two forms of contraception should include condoms, diaphragms, cervical caps, IUDs, or spermicides.
- ➔ If you are male and your partner is pregnant or capable of becoming pregnant, you should discuss with your healthcare provider the precautions you should take.

TARGRETIN Capsules can greatly increase blood levels of lipids (triglycerides and cholesterol) and these levels must be monitored and, if elevated, treated.

TARGRETIN Capsules can cause an underactive thyroid and periodic blood tests will be needed to detect this. Contact your healthcare provider if you experience symptoms such as low energy, feeling unusually cold, or always feeling tired. Medication to control the condition may be necessary.

Do not take TARGRETIN Capsules if you are allergic to this medicine.

If you have any of the following conditions, make sure you have discussed them with your healthcare provider before you start to take this medicine: You are pregnant, think you may be pregnant, or are breastfeeding; you have or previously had an inflamed pancreas (pancreatitis); you are taking gemfibrozil (Lopid®), a medication to reduce high triglyceride cholesterol (fats) levels in the blood; you are taking tamoxifen (Nolvadex®), paclitaxel (Taxol®), or atorvastatin (Lipitor®); or you are taking oral or systemic hormonal contraceptives.

Tell your healthcare provider if: you are allergic to retinoid medications; you regularly drink more than a small amount of alcohol; you are currently taking any prescription medication especially for fungal infections, bacterial infections, or seizures; or if you consume a lot of grapefruit or grapefruit juice. Tell your healthcare provider if you have or ever had: diabetes, high triglyceride (a fatty substance) levels in your blood, or gall bladder or liver disease.

Possible Complications

As an infrequent side effect of TARGRETIN Capsule treatment, pancreatitis (inflamed pancreas) may occur. If you develop persistent nausea, vomiting, abdominal or back pain while taking TARGRETIN Capsules, contact your healthcare provider immediately.

Vitamin A

Do not take more than the recommended daily dietary allowance of vitamin A (4000 to 5000 International Units).

Exposure to Sunlight

Your skin may become more sensitive to sunlight while taking this medicine. Minimize exposure to sunlight and do not use a sunlamp.

Common Side Effects

The most common side effect is an increase in blood lipids (fats in the blood). Periodic blood tests will be needed to determine blood levels of lipids, including triglycerides and cholesterol. Medication may be needed to control high fat levels in the blood.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit fda.gov/medwatch or call 1-800-FDA-1088.

Please see accompanying Prescribing Information, including Patient's Instructions for Use.